
Paper Birch



Tree Type

This tree is deciduous and will lose its foliage for the winter.

Planting

Plant in an area that is naturally moist without compacted soils. Turf grass can compete with the tree so mulching is highly recommended. Refer to back for best planting practices for your bareroot.

Watering

This tree has some drought resistance once established. Refer to watering guidelines on back for specific watering directions.

Pruning

Paper Birch may form multiple trunks so if a central leader is preferred, prune the smaller ones as soon as they sprout. Other than occasional shaping, not much pruning is needed.

Preferences

Paper Birch prefers full sun to partial shade, meaning the tree should get a minimum of 4 hours of direct, unfiltered sunlight each day. It grows well in acidic, loamy, moist, sandy, well-drained and clay soils.

Attributes

This tree grows at a medium to fast rate with height increases of anywhere from 13 to 24 inches per year but is short lived with a lifespan of 30 to 50 years. Will grow to a height of 50 to 70 ft with a spread around 35 ft at maturity. Paper Birch develops smooth white bark that curls and peels once mature and provides a bright yellow fall color. It grows in an oval shape and yields very small seeds nestled between two wings.

Wildlife Value

White-tailed deer eat a considerable amount of paper birch leaves in the fall. Snowshoe hare browse on paper birch seedlings and saplings, beavers find it a good second choice food and porcupines feed on the inner bark. Numerous cavity-nesting birds' nest in paper birch, including wood peckers, chickadees, nuthatches and swallows. Ruffed grouse eat the catskins and buds.

When newly planted trees go without enough water, growth slows to a crawl. This delays establishment and may even lead to the death of leaves, branches, roots or the whole tree.

For the most part, trees can only take up water from soil that is in direct contact with roots. Even in the best conditions, newly transplanted trees use water from a relatively small volume of soil. To make matters worse, roots of bare root, balled & burlaped, and spaded trees are cut during transplanting.

Within two to three days after spring or summer planting, the soil around the roots of trees dries enough to impede root growth. Newly transplanted trees in the Midwest benefit from daily watering for the first one to two weeks. Apply 1 to 1½ gallons of water for each inch of trunk diameter. After that, water trees every two to three days for the next two to three months and then weekly until established. The more closely you match your watering frequency to the optimum, the quicker trees become established.

Reduce watering in cool, cloudy, or wet weather if the soil is poorly drained (soil drains less than 3/4 inches per hour). Eliminate daily irrigation in poorly drained soil.

After it rains, stop watering until the rainwater drains from the soil. Stop watering in the autumn once leaves fall from trees.

Mulch reduces evaporation and conserves water. An investment in frequent watering helps insure against tree death and the cost of replanting trees.

